§ 558.360

bear caution statement as follows: Inadequate mixing (recirculation or agitation), of liquid Type B feeds has resulted in increased monensin concentration which could be fatal to goats.

(2) An approved positionally stable monensin liquid Type B feed will not be subject to the requirements for mixing directions and cautionary labeling prescribed in paragraph (f)(6)(i)(b)(1) of this section. A manufacturer may secure approval of a positionally stable liquid Type B feed by (i) either filing an NADA for the product or by establishing a master file containing data to support the stability of its product; (ii) authorizing the agency to reference and rely upon the data in the master file to support approval of a supplemental NADA to establish positional stability; and (iii) requesting No. 000986 in §510.600(c) of this chapter to file a supplemental NADA to provide for the use of its monensin Type A article in the manufacture of the liquid Type B feed specified in the appropriate master file. If the data demonstrate the stability of the liquid Type B feed described in the master file, the agency will approve the supplemental NADA. Approval of the Type B feed need not be published in the FEDERAL REGISTER because approval will not affect or alter the content of the regulation. The approval will, however, provide a basis for the individual liquid feed manufacturer to submit, and for the agency to approve, a medicated feed application under section 512(m) of the act for the liquid Type B feed. A manufacturer who seeks to market a positionally unstable monensin liquid Type B feed with mixing directions different from the standard established in paragraph (f)(6)(i)(b)(1) of this section may also follow this procedure.

(ii) [Reserved]

[40 FR 13959, Mar. 27, 1975]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting §558.355, see the List of CFR Sections Affected in the Finding Aids section of this volume.

§558.360 Morantel tartrate.

(a) Approvals. Type A medicated articles: 88 grams per pound to 000069 in $\S 510.600(c)$ of this chapter.

- (b) Related tolerances. See §556.425 of this chapter.
- (c) *Special considerations.* (1) Do not use in Type B or Type C medicated feeds containing bentonite.
- (2) Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism.
- (d) Conditions of use—(1) Amount. 0.44 to 4.4 grams of morantel tartrate per pound of feed.
- (2) Indications for use—(i) Cattle. For removal and control of mature gastro-intestinal nematode infections of cattle including stomach worms (Haemonchus spp., Ostertagia spp., Trichostrongylus spp.), worms of the small intestine (Cooperia spp., Trichostrongylus spp., Nematodirus spp.), and worms of the large intestine (Oesophagostomum radiatum).
- (ii) Goats. For removal and control of mature gastrointestinal nematode infections of goats including Haemonchus contortus, Ostertagia (Teladorsagia) circumcincta, and Trichostrongylus axei.
- (3) Limitations. Feed as a single therapeutic treatment at 0.44 gram of morantel tartrate per 100 pounds of body weight. Fresh water should be available at all times. When medicated feed is consumed, resume normal feeding. Conditions of constant worm exposure may require retreatment in 2 to 4 weeks. Do not treat cattle within 14 days of slaughter; do not treat goats within 30 days of slaughter.

[46 FR 50950, Oct. 16, 1981, as amended at 47 FR 53352, Nov. 26, 1982; 51 FR 7399, Mar. 3, 1986; 51 FR 9005, Mar. 17, 1986; 52 FR 11642, Apr. 10, 1987; 59 FR 17922, Apr. 15, 1994]

§558.363 Narasin.

- (a) Approvals. Type A medicated articles containing specified levels of narasin approved for sponsors identified in §510.600(c) of this chapter for use as in paragraph (d) of this section are as follows:
- (1) To 000986: 36, 45, 54, 72, and 90 grams per pound, paragraph (d)(1)(i) of this section.
- (2) To 000986: 36, 45, 54, 72, and 90 grams per pound, with 10, 20, 50, and 80 percent roxarsone, paragraph (d)(1)(ii) of this section.